

Table XXIV

Treatment	Reference <i>Abstracts corresponding to references can be found using the listing "RCTs by alphabetical order" or "RCTs by topic."</i>	Summary
<p>Anticoagulation (LMWH vs. Warfarin) in patients with documented thrombophilia and treated by UGFS for varices</p>	<p>Hamel-Desnos CM, Gillet JL, Desnos PR , Allaert PA. Sclerotherapy of varicose veins in patients with documented thrombophilia: a prospective controlled randomized study of 105 cases. <i>Phlebology</i>. 2009;24:176-182.</p>	<p>105 patients presenting thrombophilia: - Factor V mutation (heterozygous and homozygous) - Factor II mutation - High factor VIII and combination of 2 factors Group H: nadoparin 4000 UI simultaneously with UGFS + compression therapy 15-20 mmHg for 4 weeks Group W: warfarin 1mg per day 4 weeks after UGFS + compression therapy 15-20 mmHg for 4 weeks Results at follow-up: No episodes of symptomatic deep vein thrombosis (DVT) or pulmonary embolism (PE) occurred; no instances of DVT were revealed by US monitoring</p>

Abbreviations :

LMWH= low molecular weight heparin; US =ultrasound ; UGFS= ultrasound guided foam sclerotherapy